arXiv:1203.4972v1 [math.AG] 22 Mar 2012

APOLAR IDEAL AND NORMAL BUNDLE OF RATIONAL CURVES

ALESSANDRO BERNARDI

Abstract. As in our previous work [1] we address the problem to determine the splitting of the normal bundle of rational curves. With apolarity theory we are able to characterize some particular subvarieties in some Hilbert scheme of rational curves, defined by the splitting type of the normal bundle and the restricted tangent bundle.

1. Introduction

In this work we address the problem of classifying the rational curves $C \subset \mathbb{P}^m$ of degree n, $n \geq m$, by the splitting type of their normal bundle $N_{C;\mathbb{P}^m}$ or/and their restricted tangent bundle $T\mathbb{P}^m|_C$. This problem was addressed by several authors (see [3],[4],[6],[5],[9],[10],[11],[12],[7]). Our approach consists in a projective point of view, by considering our degree n curves as projections of the rational normal curve $C_n \subset \mathbb{P}^n$ from a linear space $L = \mathbb{P}^{k-1}$. The projected curves lie in a projective space of dimension $m = (n-k)$. We point out that we are interested in the case when C has only ordinary singularities, as in the work of Ghione and Sacchiero ([6]).

We define the scheme $H^{m,n}$ as the component of the Hilbert scheme $\mathrm{Hilb}_n \mathbb{P}^m$ of arithmetic genus zero curves of degree n in $\mathbb{P}^m$ containing the smooth curves as an open subset.

We denote with $\mathcal{N}^n_{n-k}(n_1, ..., n_{n-k-1})$ the subscheme of curves such that the splitting type of normal bundle is $(n_1, ..., n_{n-k-1})$.

One advantage in working directly on the Grassmannian $Gr(\mathbb{P}^{k-1}, \mathbb{P}^n)$ is that the number of irreducible components and the codimension of the varieties $N^n_{n-k}(n_1, ..., n_{n-k-1})$ defined as $\mathcal{N}^n_{n-k}(n_1, ..., n_{n-k-1})/SL(n-k+1)$ (parameterizing subspaces L such that the curve obtained by projecting from L has normal bundle isomorphic to $\bigoplus_{i=1}^{n-k-1} \mathcal{O}_{\mathbb{P}^1}(n_i)$) remain the same as those of $\mathcal{N}^n_{n-k}(n_1, ..., n_{n-k-1})$. Then we can study directly the basic structures of these subvarieties in the Grassmannian. In the same way we define the varieties $T^n_{n-k}(t_1, ...t_{n-k})$ with respect to the restricted tangent bundle.

In the results in [1], we have studied some Hilbert scheme of rational curves with fixed splitting type of the normal bundle and the restricted tangent bundle and we have characterized them via the individuation of some particular family of multisecant projective spaces containing L. In this work, instead, we characterize some particular subvarieties in some Hilbert

---

Date: March 1, 2013.

scheme of rational curves, defined by the splitting type of the normal bundle or the restricted tangent bundle. We obtain our main results (Theorems 3.8, 5.3) by using apolarity theory. In particular we will show that:

For $n-1 \leq 3k \leq 3(n-3)$, if the center of projection $L \cong \mathbb{P}^{k-1}$ is contained in a $\mathbb{P}^{k+1}$, $(k+2)-$secant to the rational normal curve $C_n \subset \mathbb{P}^n$, then we have:

$$N_{\pi_L(C_n);\mathbb{P}^{n-k}} \cong \mathcal{O}(n+2)^{n-k-2} \oplus \mathcal{O}(n+1+2k).$$

For $2k < n-1$, if the center of projection $L \cong \mathbb{P}^{k-1}$ is contained in a $\mathbb{P}^k$, $(k+1)-$secant to the rational normal curve $C_n \subset \mathbb{P}^n$, then we have:

$$T\mathbb{P}^{n-k}|_{\pi_k(C_n)} \cong \mathcal{O}(n+1)^{n-k-1} \oplus \mathcal{O}(n+1+k).$$

One of the main reason of interest in this work is represented by the study of the interplay between apolarity theory and the splitting type of the normal and conormal bundle (see also [1]).

2. Rational Curves of degree n in codimension k

We describe here the main steps of our approach, referring to [1] for more details. Let $C_n = \nu_n(\mathbb{P}^1) \subset \mathbb{P}^n$ be the rational normal curve which is the image of the Veronese map $\nu_n : \mathbb{P}^1 \to \mathbb{P}^n$. Let $\pi_L(C_n)$ be the rational curve obtained from C_n by projection from a k-dimensional linear subspace $L \subset \mathbb{P}^n$ on $\mathbb{P}^{n-k} \subset \mathbb{P}^n$; we will suppose that $\pi_L(C_n)$ has only ordinary singularities.

Let $J(\nu_n)$ be the Jacobian matrix of ν_n:

$$J(\nu_n) = \begin{pmatrix} ns^{n-1} & \dots & t^{n-1} & 0 \\ 0 & s^{n-1} & \dots & nt^{n-1} \end{pmatrix}.$$

We will indicate with V a 2-dimensional complex vector space, therefore we can write down in more invariant way:

(1)

$$\begin{tikzcd}
 & & 0 \arrow[d] & 0 \arrow[d] & \\
 & & \mathbb{C}^k \otimes \mathcal{O}_{C_n} \arrow[r, "\cong"] \arrow[d, "P"] & \mathbb{C}^k \otimes \mathcal{O}_{C_n} \arrow[d, "(\mathcal{N}^L_{n,k})^t"] & \\
0 \arrow[r] & V \otimes \mathcal{O}_{\mathbb{P}^1}(-n+1) \arrow[r, "J(\nu_n)"] & \operatorname{Sym}^n V \otimes \mathcal{O}_{C_n} \arrow[r, "Syz(J(\nu_n))"] \arrow[d] & \operatorname{Sym}^{n-2} V \otimes \mathcal{O}_{C_n}(2) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & V \otimes \mathcal{O}_{\mathbb{P}^1}(-n+1) \arrow[r, "J(\pi_{n-3}\circ\nu_n)"] & \frac{\operatorname{Sym}^n V}{\mathbb{C}^k} \otimes \mathcal{O}_{\pi_L(C_n)} \arrow[r] \arrow[d] & N'_{\pi_L(C_n);\mathbb{P}^{n-k}}(-n) \arrow[r] \arrow[d] & 0 \\
 & & 0 & 0. &
\end{tikzcd}$$

where the map $(\mathcal{N}^L_{n,k})^t$ is given by:

$$(\mathcal{N}^L_{n,k})^t = Syz(J(\nu_n)) \cdot \begin{pmatrix} a^1_0 & \dots & a^k_0 \\ \vdots & \ddots & \vdots \\ a^1_n & \dots & a^k_n \end{pmatrix} =$$

$$\begin{pmatrix} a^1_0t^2 - 2a^1_1ts + a^1_2s^2 & a^1_1t^2 - 2a^1_2ts + a^1_3s^2 & \dots & a^1_{n-2}t^2 - 2a^1_{n-1}ts + a^1_ns^2 \\ \vdots & \vdots & \ddots & \vdots \\ a^k_0t^2 - 2a^k_1ts + a^k_2s^2 & a^k_1t^2 - 2a^k_2ts + a^k_3s^2 & \dots & a^k_{n-2}t^2 - 2a^k_{n-1}ts + a^k_ns^2 \end{pmatrix}^t,$$

a $(n-1) \times k$ matrix. We stress out that the first row of the above diagram is SL(2) invariant, but the second one is not.

Observation 2.1. *We can observe that if $\pi_L(C_n)$ has only ordinary singularities, then the map of differential is surjective (see [6]), so ${N'}^{\vee}_{\pi_L(C_n);\mathbb{P}^{n-k}}$ is a vector bundle. We will consider only cases with ordinary singularities, so we will indicate $N'_{\pi_L(C_n);\mathbb{P}^{n-k}}$ as $N_{\pi_L(C_n);\mathbb{P}^{n-k}}$.*

We can obtain as in the case of the normal bundle of rational curves, the following exact sequence for the restricted tangent bundle :
(2)

$$\begin{tikzcd} 0 \arrow[r] & (T\mathbb{P}^{n-k}|_{\pi_L(C_n)})^{\vee}(n+1) \arrow[r] & \mathrm{Sym}^{n-1} V \otimes \mathcal{O}^n_{C_n} \arrow[r, "\mathcal{T}^L_{n,k}"] & \mathbb{C}^k \otimes \mathcal{O}_{C_n}(1) \arrow[r] & 0 \end{tikzcd},$$

where we have indicated with $T^L_{n,k}$ the $2k \times n$ matrix:

$$T^L_{n,k} = \begin{pmatrix} a^1_0 & \dots & a^1_{n-1} \\ -a^1_1 & \dots & -a^1_n \\ \vdots & \ddots & \vdots \\ a^k_0 & \dots & a^k_{n-1} \\ -a^k_1 & \dots & -a^k_n \end{pmatrix}.$$

We refer to [1] for the main results about apolarity and the Waring's Problem. Moreover we need the following results (see [8]):

Definition 2.2. Let $f \in S_n$ be a binary form of degree n. Let $L_1, ..., L_m$ be a linear forms. A representation of f as a sum:

$$f = G_1L_1^{n-g_1+1} + ... + G_sL_m^{n-g_m+1}, \tag{3}$$

where $G_i \in S_{g_i-1}$, is called a generalized additive decomposition (GAD) of f. A GAD is called normalized if no pair L_α, L_β is proportional to each other and none of the G_i is divisible by L_i. Its length is by definition $\sum_{i=1}^m g_i$.

If all $g_i = 1$ we obtain the classical additive decomposition:

$$f = c_1L_1^n + ... + c_sL_s^n, \tag{4}$$

with $c_i \in \mathbb{C}$.

The length of a binary form f is the minimum length of a GAD of f, we denote it by $l(f)$.

Lemma 2.3 ([8])**.** *Let* $\phi = \prod_{i=1}^{m}(b_i\partial_0 - a_i\partial_1)^{g_i}$ *be a prime decomposition of a nonzero form in* T_s*. Let* $L_i = a_i x_0 + b_i x_1$*. Then a form* $f \in S_n$ *with* $n \geq s$ *has a GAD as 3 if and only if* ϕ *is apolar to* f*. If all roots of* ϕ *are simple, then this is an additive decomposition.*

Lemma 2.4 ([8])**.** *Let* $n = 2t$ *or* $n = 2t+1$*, let* $f \in S_n$*. Then* $l(f) \leq t+1$*. If* $I_f = Ann(f)$ *is the ideal of forms apolar to* f*, then* $l(f)$ *equals the order* d *(i.e. the initial degree) of the graded ideal* $I_f = (I_f)_d + (I_f)_{d+1} + \ldots$*.*

Lemma 2.5 (Jordan's Lemma)**.** *Suppose that the linear forms* $L_i, i = 1, \ldots, m$ *are not proportional to each other and:*

$$0 = G_1 L_1^{n-g_1+1} + \ldots + G_s L_m^{n-g_m+1},$$

with $\sum_{i=1}^{m} g_i \leq n+1$*. Then* $G_i = 0$ *for every* i*.*

Proposition 2.6 (Uniqueness of GAD)**.** *Suppose* $n = 2t$ *or* $n = 2t+1$*. Let:*

$$f = G_1 L_1^{n-g_1+1} + \ldots + G_m L_m^{n-g_m+1},$$

be a normalized GAD of $f \in S_n$ *of length* $s = \sum_{i=1}^{m} g_i \leq t+1$*. Then* f *has no other GAD of length* $\leq n+1-s$ *and* $l(f) = s$*. In particular if* $s \leq t$ *or if* $s = t+1, n = 2t+1$ *(equivalently* $2s \leq n+1$*), then the above is the unique normalized GAD of* f *having length* $\leq t+1$*.*

Definition 2.7. Let $f \in S_n$ and let $2l(f) \leq n+1$. Then the unique normalized GAD of length $s = l(f)$ is called the canonical form of f.

Theorem 2.8 (Sylvester)**.**

i) *For odd* $n = 2t+1$*, the general* $f \in S_n$ *has a unique decomposition as a sum of* $t+1$ $n-$*th powers of linear forms.*

ii) *For even* $n = 2d$*, the general* $f \in S_n$ *has infinitely many decompositions as a sum of* $t+1$ $n-$*th powers of linear forms.*

Theorem 2.9. *Let* $n = 2t$ *or* $2t+1$*, let* $f \in S_n$*.*

i) *Let* $s = \operatorname{rank}\operatorname{Cat}_f(n-t, t; 2)$*. Then* $l(f) = s$*. If* $2s \leq n+1$*, then* f *has a unique generalized additive decomposition of length* s*. and no other GADs of length* $\leq t+1$*.*

ii) *For every pair of integers* s, e *with* $1 \leq s \leq e \leq n-e+1$*, if* $l(f) = s$*, then* $l(f) = s = \operatorname{rank}\operatorname{Cat}_f(n-e, e; 2)$*.*

Theorem 2.10 ([8])**.** *Let* f *be a binary form of degree* $n = 2t$ *or* $2t+1$ *and* $I_f = Ann(f)$ *be the ideal of forms apolar to* f*. Let* $A_f = S/I_f$ *be the associated Gorenstein Artin algebra. Let* $s = \max\{\dim(A_f)_i\}$*. Then:*

i. $s = l(f)$ *and the Hilbert function of* A_f *satisfies:*

$$H(A_f) = (1, 2, \ldots, s-1, \overset{s-1}{s}, s, \ldots, \overset{n-s+1}{s}, s-1, \ldots, 2, 1);$$

ii. *Suppose* $2s \leq n+1$*. Then* $\dim I_s = 1$*,* $I_s =< \alpha >$ *and for every integer* v *with* $s \leq v \leq n-s+1$ *one has* $I_v = S_{v-s} \circ \alpha$*;*

iii. *The apolar ideal I_f is generated by two homogeneous polynomials $\alpha \in (I_f)_s$ and $\beta \in (I_f)_{n+2-s}$.*
Equivalently the ring A_f is a complete intersection of generator degrees $s, n+2-s$. The two polynomials above have no common zeros.

In this contest our principal tool will be the Apolarity Lemma (see [8]):

Lemma 2.11 (Apolarity Lemma). *Let $p_1, ..., p_s \in \mathbb{P}^r$, let $L_i = L_{p_i}$, let $P = \{[p_1], ..., [p_s]\} \subset \mathbb{P}^r$ and let $\mathcal{I}_P$ be the homogeneous ideal in T of polynomials vanishing on P. Then:*

i) *For every $\phi \in R_e$:*
$$\phi \circ (L_1^{[N]} + ... + L_s^{[N]}) = \phi(p_1)L_1^{[N-E]} + ... + \phi(p_s)L_s^{[N-E]}.$$
ii) *With respect to the contraction paring $T_n \times S_n \to \mathbb{C}$ one has:*
$$((\mathcal{I}_P)_n)^{\perp} =< L_1^{[N]}, ..., L_s^{[N]} > .$$
iii) *The points $[p_1], ..., [p_s] \subset \mathbb{P}^r$ impose independent conditions on the linear system $|\mathcal{O}_{\mathbb{P}^n}(j)|$ if and only if $L_1^{[N]}, ..., L_s^{[N]}$ are linearly independent.*
iv) *Suppose $s \leq \dim_{\mathbb{C}} T_{n-e}$ and the linear forms $L_1^{[N]}, ..., L_s^{[N]}$ have the property that the corresponding set P imposes independent conditions on the linear system $|\mathcal{O}_{\mathbb{P}^r}(n-e)|$. Let $f = L_1^{[N]} + ... + L_s^{[N]}$. Then we have for the apolar forms to f of degree e the equality:*
$$Ann(f)_e = (\mathcal{I}_P)_e.$$

2.1. Normal Bundle of Rational Curves in $\mathbb{P}^{n-1}$.

It is easy to prove the following result:

Theorem 2.12. *Let $C_n \subset \mathbb{P}^n$ be the rational normal curve of degree n. For any $p \in \mathbb{P}^n$, the rational curve $\pi_p(C_n) \subset \mathbb{P}^{n-1}$ has normal bundle $N_{\pi_p(C_n);\mathbb{P}^{n-1}} = \mathcal{O}(n+2)^{n-4} \oplus \mathcal{O}(n+3)^2$ if and only if p is not on a secant (or tangent) line to C_n. This is equivalent to saying that $\pi_p(C_n)$ is smooth.*

2.2. Normal Bundle of Rational Curves in $\mathbb{P}^{n-2}$.

By diagram (1) in case of codimension 2 since the map $\mathcal{O}_{C_n}(2)^{n-1} \to N_{\pi_L(C_n);\mathbb{P}^{n-2}}(-n)$ is surjective, it follows that $2 \leq n'_0 \leq ... \leq n'_{n-4}$ and $n'_0 + ... + n'_{n-4} = 2n-2$, where $N_{\pi_L(C_n);\mathbb{P}^{n-2}}(-n) = \mathcal{O}(n'_0) \oplus ... \oplus \mathcal{O}(n'_{n-4})$. But we can write $n'_i = n''_i + 2$, so $n''_0 + ... + n''_{n-4} = 4$.

Therefore we have $n'_0 = ... = n'_{n-8} = 2$, we can write:
$$N_{\pi_2(C_n);\mathbb{P}^{n-2}} = \mathcal{O}(n+2)^{n-7} \oplus \mathcal{F};$$
where $\mathcal{F}$ is a rank 4 vector bundle on $\mathbb{P}^1$. Hence we must study the splitting of $\mathcal{F}$, if we indicate with $\mathcal{F} = \mathcal{O}(f_0) \oplus ... \oplus \mathcal{O}(f_3)$, where $f_0 + ... + f_3 = 4n+14$. Therefore it is one of the following cases:

(1) $\mathcal{F} = \mathcal{O}(n+5)^4 =: \mathcal{F}_1$;

(2) $\mathcal{F} = \mathcal{O}(n+4) \oplus \mathcal{O}(n+5)^2 \oplus \mathcal{O}(n+6) =: \mathcal{F}_2$;
(3) $\mathcal{F} = \mathcal{O}(n+4)^2 \oplus \mathcal{O}(n+6)^2 =: \mathcal{F}_3$;
(4) $\mathcal{F} = \mathcal{O}(n+4)^2 \oplus \mathcal{O}(n+5) \oplus \mathcal{O}(n+7) =: \mathcal{F}_4$;
(5) $\mathcal{F} = \mathcal{O}(n+4)^3 \oplus \mathcal{O}(n+8) =: \mathcal{F}_5$;

no other case can occur if the projection has only ordinary singularities by Observation.2.1.

Since the codimension is 2, by dualizing the last exact column of (1) and tensorizing with $\mathcal{O}_{\mathbb{P}^1}(2)$, we get:

$$\begin{tikzcd} 0 \arrow[r] & N^{\vee}_{\pi_2(C_n);\mathbb{P}^{n-2}}(n+2) \arrow[r] & \mathcal{O}^{n-1}_{C_n} \arrow[r, "\mathcal{N}^L_{n,2}"] & \mathcal{O}^2_{C_n}(2) \arrow[r] & 0, \end{tikzcd}$$

and we have $deg(N^{\vee}_{\pi_L(C_n);\mathbb{P}^{n-2}}(n+2)) = -2$. But $N^{\vee}_{\pi_L(C_n);\mathbb{P}^{n-2}}(n+2) = \mathcal{O}^{n-7} \oplus \mathcal{F}^{\vee}(n+2)$, so $h^0(N^{\vee}_{\pi_L(C_n);\mathbb{P}^{n-2}}(n+2)) = n - 7 + h^0(\mathcal{F}^{\vee}(n+2))$. Therefore we have that $0 \leq h^0(\mathcal{F}^{\vee}(n+2)) \leq 3$, where:

a) $h^0(\mathcal{F}^{\vee}(n+2)) = 0 \Leftrightarrow \mathcal{F} = \mathcal{F}_1$;
b) $h^0(\mathcal{F}^{\vee}(n+2)) = 1 \Leftrightarrow \mathcal{F} = \mathcal{F}_2$;
c) $h^0(\mathcal{F}^{\vee}(n+2)) = 2 \Leftrightarrow \mathcal{F} = \mathcal{F}_3$ *or* $\mathcal{F}_4$;
d) $h^0(\mathcal{F}^{\vee}(n+2)) = 3 \Leftrightarrow \mathcal{F} = \mathcal{F}_5$.

So we have the following cases:

A) $\mathrm{rank}(N^L_{n,2}) = 6 \Leftrightarrow h^0(\mathcal{F}^{\vee}(2)) = 0 \Leftrightarrow \mathcal{F} = \mathcal{F}_1$;
B) $\mathrm{rank}(N^L_{n,2}) = 5 \Leftrightarrow h^0(\mathcal{F}^{\vee}(2)) = 1 \Leftrightarrow \mathcal{F} = \mathcal{F}_2$;
C) $\mathrm{rank}(N^L_{n,2}) = 4 \Leftrightarrow h^0(\mathcal{F}^{\vee}(2)) = 2 \Leftrightarrow \mathcal{F} = \mathcal{F}_3$ *or* $\mathcal{F}_4$;
D) $\mathrm{rank}(N^L_{n,2}) = 3 \Leftrightarrow h^0(\mathcal{F}^{\vee}(2)) = 3 \Leftrightarrow \mathcal{F} = \mathcal{F}_5$.

If we consider the following exact sequence:

$$\begin{tikzcd} 0 \arrow[r] & N^{\vee}_{\pi_L(C_n);\mathbb{P}^{n-2}}(n+3) \arrow[r] & \mathcal{O}^{n-1}_{C_n}(1) \arrow[r, "\mathcal{N}^L_{n,2}"] & \mathcal{O}^2_{C_n}(3) \arrow[r] & 0, \end{tikzcd}$$

we have that:

C1) $\mathrm{rank}(N^L_{n,2}(1)) = 3 \Leftrightarrow h^0(\mathcal{F}^{\vee}(3)) = 8 \Leftrightarrow \mathcal{F} = \mathcal{F}_3$;
C2) $\mathrm{rank}(N^L_{n,2}(1)) \leq 4 \Leftrightarrow h^0(\mathcal{F}^{\vee}(3)) = 7 \Leftrightarrow \mathcal{F} = \mathcal{F}_4$.

Proposition 2.13. *If there exist two points $q_1, q_2 \in L$, each of them belonging to a different $3-$secant $\mathbb{P}^2$, then:*

$$\ker(N^L_{n,2}) = (\mathcal{I}_{D_1} \cap \mathcal{I}_{D_2})_{n-2} = (\mathcal{I}_{D_1 \cup D_2})_{n-2},$$

where D_i is the set of points in $\mathbb{P}^1$ which corresponds to the linear forms in the additive decomposition of f_i (the binary form corresponding to q_i).

2.2.1. *Case* $\mathrm{rank}\, N^L_{n,2} = 3$.

Lemma 2.14. *The splitting type of the normal bundle $N_{\pi_L(C_n),\mathbb{P}^{n-2}}$ is:*

$$((n+2)^{n-4}, n+6)$$

if and only if $\mathrm{rank}(N^L_{n,2}) = 3$.

Theorem 2.15 (Case Rank 3)**.** *If the projection line L belongs to some $3-$secant $\mathbb{P}^2$, but it is not a secant line, then the splitting type of the normal bundle $N_{\pi_L(C_n),\mathbb{P}^{n-2}}$ is:*

$$((n+2)^{n-4}, n+6).$$

Proof. If L belongs to a $3-$secant $\mathbb{P}^2$, then there exist two points $q_1, q_2 \in L$ and their corresponding binary forms f_1, f_2 of degree n such that $Ann(f_1) = (\alpha, \beta_1)$ and $Ann(f_2) = (\alpha, \beta_2)$ where α has only simple roots, $\deg(\alpha) = 3$ and $\beta_1 \neq \beta_2$, $\deg(\beta_1) = \deg(\beta_2) = n-1$. So dim $<\alpha>_{n-2} = n-4$ and $<\alpha, \beta_1>_{n-2} = <\alpha, \beta_2>_{n-2} = <\alpha>_{n-2}$. Therefore $\text{rank}(N^L_{n,2}) = n-1-\dim(<\alpha,\beta_1>_{n-2} \cap <\alpha,\beta_2>_{n-2}) = 3$.

□

Corollary 2.16. *The variety of lines L such that π_L gives a rational curve of degree n in $\mathbb{P}^{n-2}$ for which the splitting type of the normal bundle $N_{\pi_L(C_n),\mathbb{P}^{n-2}}$ is:*

$$((n+2)^{n-4}, n+6)$$

has an irreducible subvariety of codimension $(2n-7)$ in $Gr(\mathbb{P}^1, \mathbb{P}^n)$, that is formed by the lines belonging to some $3-$secant $\mathbb{P}^2$, but which are not secant lines.

2.2.2. *Case* rank $N^L_{n,2} = 4$.

Lemma 2.17. *The codimension in $Gr(\mathbb{P}^1, \mathbb{P}^n)$ of the variety of all lines in $\mathbb{P}^n$ belonging to some 4-secant $\mathbb{P}^3$ to the rational normal curve in $\mathbb{P}^n$ is $2n-10$*

Proof. In fact we can consider the incidence variety $I = \{(L,\pi) : L \in Gr(\mathbb{P}^1, \mathbb{P}^n), \pi \in S, L \subset S\}$ where S is the set of all 4-secant $\mathbb{P}^3$. In the usual way we can compute the codimension of the image of this incidence variety in $Gr(\mathbb{P}^1, \mathbb{P}^n)$. That calculation is effective thanks to the result of Chiantini and Ciliberto on the non-defectivity of the Grassmannians of secant varieties of curves (see [2]). □

Lemma 2.18. *If the splitting type of the normal bundle $N_{\pi_L(C_n),\mathbb{P}^{n-2}}$ is:*

$$((n+2)^{n-5}, (n+4)^2),$$

then $\text{rank}(N^L_{n,2}) = 4$. *Moreover the variety which parameterizes the lines giving the above splitting has codimension $2(n-5)$ in $Gr(\mathbb{P}^1, \mathbb{P}^n)$.*

Theorem 2.19 (Case Rank 4)**.** *If the projection line L belongs to some $4-$secant $\mathbb{P}^3$, but it does not belong to some $3-$secant $\mathbb{P}^2$, then the splitting type of the normal bundle $N_{\pi_L(C_n),\mathbb{P}^{n-2}}$ is:*

$$((n+2)^{n-5}, (n+4)^2).$$

Proof. If L belongs to a $4-$secant $\mathbb{P}^3$, then there exist two points $q_1, q_2 \in L$ and their corresponding binary forms f_1, f_2 of degree n such that $Ann(f_1) = (\alpha, \beta_1)$ and $Ann(f_2) = (\alpha, \beta_2)$ with α has only simple roots and $\deg(\alpha) = 4$ and $\beta_1 \neq \beta_2$, $\deg(\beta_1) = \deg(\beta_2) = n-2$. So dim $<\alpha>_{n-2} = n-5$ and

$\dim(<\beta_1>_{n-2}\cap<\beta_2>_{n-2})=0$ otherwise $\beta_1=\beta_2$ and $q_1=q_2$, but this is impossible. Therefore $\operatorname{rank}(N^L_{n,2})=n-1-\dim(<\alpha,\beta_1>_{n-2}\cap<\alpha,\beta_2>_{n-2})=4$.

□

Corollary 2.20. *The variety of lines $L\subset\mathbb{P}^n$ such that $\pi_L(C_n)\subset\mathbb{P}^{n-2}$ has the splitting type of the normal bundle $N_{\pi_L(C_n),\mathbb{P}^{n-2}}$:*

$$((n+2)^{n-5},(n+4)^2)$$

has an irreducible component formed by the lines belonging to some $4-$secant $\mathbb{P}^3$, but not contained in any $3-$secant $\mathbb{P}^2$.

2.2.3. *Case* $\operatorname{rank}N^L_{n,2}=5$.

Lemma 2.21. *The splitting type of the normal bundle $N_{\pi_L(C_n),\mathbb{P}^{n-2}}$ is:*

$$((n+2)^{n-6},(n+3)^2,n+4)$$

if and only if $\operatorname{rank}(N^L_{n,2})=5$.

Observation 2.22. *If L belongs to a $5-$secant $\mathbb{P}^4$, then there exist two points $q_1,q_2\in L$ and their corresponding binary forms f_1,f_2 of degree n such that $Ann(f_1)=(\alpha,\beta_1)$ and $Ann(f_2)=(\alpha,\beta_2)$ with α has only simple roots and* $\deg(\alpha)=5$ *and* $\beta_1\neq\beta_2$, $\deg(\beta_1)=\deg(\beta_2)=n-3$. *So* $\dim<\alpha>_{n-2}=n-6$, *so* $\operatorname{rank}(N^L_{n,2})=n-1-\dim(<\alpha,\beta_1>_{n-2}\cap<\alpha,\beta_2>_{n-2})\leq 5$.

Observation 2.23. *If* $\operatorname{rank}(N^L_{n,2})=5$, *then or $L\subset\mathbb{P}^4$ which is $5-$secant to C_n or $L\subset\mathbb{P}^k$ which is $(k+1)-$secant. In the second case there exist two points $q_1,q_2\in L$ such that $Ann(f_1)=(\alpha,\beta_1)$ and $Ann(f_2)=(\alpha,\beta_2)$ with α has only simple roots and* $\deg(\alpha)=k+1$ *and* $\beta_1\neq\beta_2$, $\deg(\beta_1)=\deg(\beta_2)=n-k+1$.

2.3. **Normal Bundle of Rational Curves in $\mathbb{P}^{n-3}$.**

2.3.1. *Case* $\operatorname{rank}N^L_{n,3}=6,5$.

Observation 2.24. *If $L\cong\mathbb{P}^2$ is contained in a $6-$secant $\mathbb{P}^5$ to C_n, then there exist three binary forms of degree n correspond to three points $q_1,q_2,q_3\in L$ such that $Ann(f_1)=(\alpha,\beta_1),Ann(f_2)=(\alpha,\beta_2)$ and $Ann(f_3)=(\alpha,\beta_3)$ where α has only simple roots,* $\deg(\alpha)=6$ *and* $\beta_1\neq\beta_2\neq\beta_3$, $\deg(\beta_1)=\deg(\beta_2)=\deg(\beta_3)=n-4$. *So* $\dim<\alpha>_{n-2}=n-7$, *so* $\operatorname{rank}(N^L_{n,2})=n-1-\dim(<\alpha,\beta_1>_{n-2}\cap<\alpha,\beta_2>_{n-2})\leq 6$.

Observation 2.25. *If $L\cong\mathbb{P}^2$ is contained in a a $5-$secant $\mathbb{P}^4$, then there exist three binary forms of degree n correspond to three points $q_1,q_2,q_3\in L$ such that $Ann(f_1)=(\alpha,\beta_1),Ann(f_2)=(\alpha,\beta_2)$ and $Ann(f_3)=(\alpha,\beta_3)$ where α has only simple roots,* $\deg(\alpha)=5$ *and* $\beta_1\neq\beta_2\neq\beta_3$, $\deg(\beta_1)=\deg(\beta_2)=\deg(\beta_3)=n-3$. *So* $\dim<\alpha>_{n-2}=n-6$, *and* $\operatorname{rank}(N^L_{n,2})=n-1-\dim(<\alpha,\beta_1>_{n-2}\cap<\alpha,\beta_2>_{n-2})\leq 5$.

2.3.2. *Case* $\operatorname{rank} N^L_{n,3} = 4$.

Proposition 2.26. *If* $L \cong \mathbb{P}^2$ *is contained in a* $4-$*secant* $\mathbb{P}^3$*, but it is not a* $3-$*secant* $\mathbb{P}^2$*, then the splitting type of the normal bundle* $N_{\pi_L(C_n),\mathbb{P}^{n-3}}$ *is:*

$$((n+2)^{n-5}, n+8).$$

Such L's form an irreducible component of codimension $(3n-13)$ *of the varieties of projection planes which give the splitting type above .*

Proof. If $L \cong \mathbb{P}^2$, as centre of projection, belongs to a $4-$secant $\mathbb{P}^3$, then there exist three binary forms of degree n correspond to three points $q_1, q_2, q_3 \in L$ such that $Ann(f_1) = (\alpha, \beta_1)$, $Ann(f_2) = (\alpha, \beta_2)$ and $Ann(f_3) = (\alpha, \beta_3)$ with α has only simple roots and $\deg(\alpha) = 4$ and $\beta_1 \neq \beta_2 \neq \beta_3$, $\deg(\beta_1) = \deg(\beta_2) = \deg(\beta_3) = n-2$. So $\dim <\alpha>_{n-2} = n-5$, so $\operatorname{rank}(N^L_{n,3}) = n-1-\dim(<\alpha,\beta_1>_{n-2} \cap <\alpha,\beta_2>_{n-2} \cap <\alpha,\beta_3>_{n-2}) = 4$, otherwise $\beta_1 = \beta_2 = \beta_3$, but this is impossible. ☐

3. Normal Bundle of Rational Curves in $\mathbb{P}^{n-k}$, for $\frac{n-1}{3} \leq k \leq n-3$

Observation 3.1. *We always have:*

$$k \leq \operatorname{rank} N^L_{n,k} = n-1-h^0(N^\vee_{\pi_L(C_n),\mathbb{P}^{n-k}}(n+2)) \leq n-1.$$

Proposition 3.2. $N_{\pi_L(C_n);\mathbb{P}^{n-k}} \cong \mathcal{O}(n+2)^{n-1-\operatorname{rank}(N^L_{n,k})} \oplus \mathcal{F}$*, where* $\mathcal{F}$ *is a vector bundle on* $\mathbb{P}^1$ *of* $\operatorname{rank}(N^L_{n,k}) - k$ *on* $\mathbb{P}^1$ *and* $\deg(\mathcal{F}^\vee(n+2)) = -2k$.
We have $\mathcal{F} \cong \bigoplus_{i=0}^{\operatorname{rank}(N^L_{n,k})-k} \mathcal{O}(l_i)$ *with* $l_i \geq n+3$.
If $2(n-k) \geq 2k$ *we have two possibilities:*

1. $N_{\pi_L(C_n);\mathbb{P}^{n-k}} \cong \mathcal{O}(n+2)^{r-1} \oplus \mathcal{O}(n+3)^{2(n-k-r)-2k} \oplus \mathcal{F}'$ *with* $\operatorname{rank}(\mathcal{F}') = n-r-2(n-k-r)+k$ *and* $\deg(\mathcal{F}'^\vee(n+2)) = -2k+2(2(n-k-r)-2k)$ *if and only if* $\operatorname{rank}(N^L_{n,k}) = n-r$ *and* $2(n-k-r) \geq 2k$ *for* $1 \leq r \leq n-k-2$;
2. $N_{\pi_L(C_n);\mathbb{P}^{n-k}} \cong \mathcal{O}(n+2)^{n-k-2} \oplus \mathcal{O}(n+2+2k)$ *if and only if* $\operatorname{rank}(N^L_{n,k}) = k+1$.

However the last one is true also for $2n-2k < 2k$.

We can rephrase the above proposition as:

Proposition 3.3. *If* $2(n-k) \geq 2k$ *we have two possibilities:*

1. $\pi_L(C_n) \in N^n_{n-k}((n+2)^{r-1}, (n+3)^{2(n-k-r)-k}, spt(\mathcal{F}'))$*, where* $spt(\mathcal{F}')$ *is the splitting type of* $\mathcal{F}'$ *with* $\operatorname{rank}(\mathcal{F}') = n-r-2(n-k-r)$ *and* $\deg(\mathcal{F}'^\vee(n+2)) = -2k+2(2(n-k-r)-2k)$ *if and only if* $L \in V(N^L_{n,k})^{n-r}$ *and* $2(n-k-r) \geq 2k$ *for* $1 \leq r \leq n-k-2$;
2. $\pi_L(C_n) \in N^n_{n-k}((n+2)^{n-k-1}, (n+2+k))$ *if and only if* $L \in V(N^L_{n,k})^{k+1}$.

However the last one is true also for $2n-2k < 2k$.

Lemma 3.4. $\operatorname{rank} N^L_{n,k} \leq n-2$ *if and only if the forms* f_i *of degree* n *corresponding to the points* p_i *generating* L *can be represented by the similar GAD, i.e. :*

$$f_i = G_{i_1} L_1^{n-g_1+1} + \ldots + G_{i_m} L_m^{n-g_m+1}.$$

Proof. $\Rightarrow$ If $\operatorname{rank} N^L_{n,k} \leq n-2$, then there exists at least an element $\phi \in T_{n-2}$ such that for all forms f_i corresponding to the points p_i generating L we have $\phi \circ f_i = 0$. So we can consider the primary decomposition of $\phi = \prod_{i=1}^m (\phi_i)^{g_i}$, with $\phi_i \in T_1$ and $\sum_i g_i = n-2$, so every f_i can be represented by the similar GAD, i.e. :

$$f_i = G_{i_1} L_1^{n-g_1+1} + \ldots + G_{i_m} L_m^{n-g_m+1},$$

where $(L_j)^\perp = \phi_j$ for all $j = 1, .., m$ and $G_{i_j} \in S_{g_j-1}$ for all $i = 1, \ldots, k$ and $j = 1, \ldots, m$.

$\Leftarrow$ On the other hand if every f_i can be represented by the similar GAD, i.e. :

$$f_i = G_{i_1} L_1^{n-g_1+1} + \ldots + G_{i_m} L_m^{n-g_m+1},$$

then we can consider $\phi = \prod_{i=1}^m ((L_i)^\perp)^{g_i}$. By definition of GAD representation we have $\phi \circ f_i = 0$ for all $i = 1, \ldots, k$, so $\phi \in \ker N^L_{n,k}$ and $\operatorname{rank} N^L_{n,k} \leq n-2$.

□

Observation 3.5. *In particular we can observe that if* L *belong to a* $(n-2)-$*secant* $\mathbb{P}^{n-3}$ *generated by* $q_1, \ldots, q_{n-2}$*, then there exists an element* $\phi \in H^0(\mathcal{O}^{n-1}_{\pi_L(C_n)}) \cong S^{n-2}V^\vee = T_{n-2}$ *such that* $\phi \in \ker(N^L_{n,k}) = \bigcap_i \ker(Cat_{f_i}(2, n-2))$*, in fact we can take* $\phi = \prod_{i=1}^{n-2} L^\perp_{q_i}$*.*

We can compute the codimension of the variety of every $\mathbb{P}^{k-1}$ *which belongs to some* $(n-2)-$*secant* $\mathbb{P}^{n-3}$ *by constructing an incidence variety:*

$$I_S = \{(L, \pi) : L \in Gr(\mathbb{P}^{k-1}, \mathbb{P}^n), \pi \in S, L \subset S\},$$

where S *is the set of all* $(n-2)$*-secant* $\mathbb{P}^{n-3}$ *to* C_n*. In the usual way we can compute the codimension of the image of this incidence variety in* $Gr(\mathbb{P}^{k-1}, \mathbb{P}^n)$*. We will indicate with* ϕ_1 *and* ϕ_2 *the natural projections:*

$$\begin{tikzcd}
 & I_S \arrow[dl, "\phi_1"'] \arrow[dr, "\phi_2"] & \\
Gr(\mathbb{P}^{k-1}, \mathbb{P}^n) & & S,
\end{tikzcd}$$

so the codimension in $Gr(\mathbb{P}^{k-1}, \mathbb{P}^n)$ *of* $\phi_1(I_S)$ *is equal to* $\dim Gr(\mathbb{P}^{k-1}, \mathbb{P}^n) - \dim S - \dim \phi_2^{-1}(S) = k(n+1-k) - n + 2 - k(n-2-k)$*. The above calculation is effective thanks to the result of Chiantini and Ciliberto on the non-defectivity of the Grassmannians of secant varieties of curves (see* [2]*). We have that this variety has codimension* $3k-n+2$ *which is the codimension expected as determinantal variety.*

In general we can prove:

Lemma 3.6. *If the centre of projection* $L \cong \mathbb{P}^{k-1}$ *belongs to some* $(n-1-r)-$*secant* $\mathbb{P}^{n-r-2}$ *to the rational normal curve* C_n *in* $\mathbb{P}^n$*, then we have* $\operatorname{rank} N^L_{n,k} \leq n-r$ *for* $1 \leq r < n-k-1$.

Proof. If $L \cong \mathbb{P}^{k-1}$ belongs to some $(n-r-1)-$secant $\mathbb{P}^{n-r-2}$, then there exist k points $p_1, ..., p_k \in L$ which generate L and the corresponding binary forms f_i are generated by two forms $Ann(f_i) = (\alpha, \beta_i)$ where $\deg(\alpha) = n-r-1$, α has only simple roots and $\deg(\beta_i) = r+3$ without common zeros with α. We have $\dim(\alpha)_{n-2} = r$ and $\dim \bigcap_i (\alpha, \beta_i)_{n-2} \geq r$, so $\operatorname{rank} N^L_{n,k} = n-1-\dim \bigcap_i (\alpha, \beta_i)_{n-2} \leq n-r-1$. □

Lemma 3.7. *The codimension in* $Gr(\mathbb{P}^{k-1}, \mathbb{P}^n)$ *of the variety of all* $L \cong \mathbb{P}^{k-1}$ *in* $\mathbb{P}^n$ *belonging to some* $(n-r-1)$*-secant* $\mathbb{P}^{n-r-2}$ *to the rational normal curve in* $\mathbb{P}^n$ *is* $2k+kr-n+r+1$.

Proof. Infact we can consider the incidence variety $I_S = \{(L, \pi) : L \in Gr(\mathbb{P}^{k-1}, \mathbb{P}^n), \pi \in S, L \subset S\}$ where S is the set of all $(n-r-1)$-secant $\mathbb{P}^{n-r-2}$ to the rational normal curve in $\mathbb{P}^n$. In the usual way we can compute the codimension of the image of this incidence variety in $Gr(\mathbb{P}^{k-1}, \mathbb{P}^n)$. The above calculation is effective thanks to the result of Chiantini and Ciliberto on the non-defectivity of the Grassmannians of secant varieties of curves (see [2]). □

In particular for $r = n-k-1$ we have:

Theorem 3.8. *If the centre of projection* $L \cong \mathbb{P}^{k-1}$ *lies in to some* $(k+2)-$*secant* $\mathbb{P}^{k+1}$ *to the rational normal curve* C_n *in* $\mathbb{P}^n$*, then we have:*

$$N_{\pi_L(C_n);\mathbb{P}^{n-k}} \cong \mathcal{O}(n+2)^{n-k-2} \oplus \mathcal{O}(n+1+2k).$$

Observation 3.9. $k+1 \leq \operatorname{rank} N^L_{n,k} = n-1-h^0(N^\vee_{\pi_L(C_n),\mathbb{P}^{n-k}}(n+2)) \leq 3k$

Proposition 3.10. $N_{\pi_L(C_n);\mathbb{P}^{n-k}} \cong \mathcal{O}(n+2)^{n-1-\operatorname{rank}(N^L_{n,k})} \oplus \mathcal{F}$*, with* $\mathcal{F}$ *a vector bundle of rank* $\operatorname{rank}(N^L_{n,k}) - k$ *on* $\mathbb{P}^1$ *and* $\deg(\mathcal{F}^\vee(n+2)) = -2k$ *such that* $\mathcal{F} \cong \bigoplus_i^{\operatorname{rank}(N^L_{n,k})-k} \mathcal{O}(l_i)$ *with* $l_i \geq n+3$.

In this case we have three possibilities:

1. $\mathcal{F} \cong \mathcal{O}(n+3)^{2k}$ *if and only if* $\operatorname{rank}(N^L_{n,2}) = 3k$*;*
2. $\mathcal{F} \cong \mathcal{O}(n+3)^{2k-2} \oplus \mathcal{O}(n+4)$ *if and only if* $\operatorname{rank}(N^L_{n,k}) = 3k-1$*;*
3. $\mathcal{F} \cong \mathcal{O}(n+3)^{2k-2r} \oplus \mathcal{F}'$ *with* $\operatorname{rank}(\mathcal{F}') = r$ *and* $\deg(\mathcal{F}'^\vee(n+2)) = -2k$ *if and only if* $\operatorname{rank}(N^L_{n,k}) = 3k-r$ *with* $1 < r \leq 2k-1$*;*
4. $\mathcal{F} \cong \mathcal{O}(n+2+2k)$ *if and only if* $\operatorname{rank}(N^L_{n,k}) = k+1$.

4. NORMAL BUNDLE OF RATIONAL CURVES IN $\mathbb{P}^{n-k}$, FOR $k < \frac{n-1}{3}$

Observation 4.1. *If* $L \cong \mathbb{P}^{k-1}$*, as centre of projection, belongs to a* $(k+1)-$*secant* $\mathbb{P}^k$*, then there exist* k *binary forms of degree* n *correspond to points*

$q_1, .., q_k \in L$ such that $Ann(f_1) = (\alpha, \beta_1), ..., Ann(f_k) = (\alpha, \beta_r)$ with α has only simple roots and $\deg(\alpha) = k + 1$ and $\beta_1 \neq ... \neq \beta_k$, $\deg(\beta_1) = ... = \deg(\beta_k) = n - k + 1$, where f_i is the binary form correspond to the point q_i. So $\dim < \alpha >_{n-2} = n - k - 2$ and $\dim < \beta_i >_{n-2} = k$, therefore $n - k - 1 \leq \mathrm{rank}(N^L_{n,k}) = n - 1 - \dim(< \alpha, \beta_1 >_{n-2} \cap ... \cap < \alpha, \beta_k >_{n-2}) \leq k + 1$.

Observation 4.2. *We can observe that if L belongs to a $(n-2)-$secant $\mathbb{P}^{n-3}$ generated by $q_1, ..., q_{n-1}$, then there exists an element $\phi \in H^0(\mathcal{O}^n_{\pi_L(C_n)}) \cong S^{n-1}V^\vee = T_{n-1}$ such that $\phi \in \ker(N^L_{n,k}) = \bigcap_i \ker(Cat_{f_i}(2, n-2))$, in fact we can always take $\phi = \prod_{i=1}^n L^\perp_{q_i}$, since $\dim \ker N^L_{n,k} \geq 1$.*

Unfortunately this condition is empty for $3k < n - 2$, in fact we can compute the codimension of the variety of every $\mathbb{P}^{k-1}$ which belong to some (n-2)-secant $\mathbb{P}^{n-3}$ constructing an incidence variety:

$$I_S = \{(L, \pi) : L \in Gr(\mathbb{P}^{k-1}, \mathbb{P}^n), \pi \in S, L \subset S\},$$

where S is the set of all (n-2)-secant $\mathbb{P}^{n-3}$ to C_n. In the usual way we can compute the codimension of the image of this incidence variety in $\mathbb{G}r(k-1, n)$. We will indicated with ϕ_1 and ϕ_2 the natural projections:

$$\begin{tikzcd}
& I_S \arrow[dl, "\phi_1"'] \arrow[dr, "\phi_2"] & \\
Gr(\mathbb{P}^{k-1}, \mathbb{P}^n) & & S,
\end{tikzcd}$$

so the codimension in $Gr(\mathbb{P}^{k-1}, \mathbb{P}^n)$ of $\phi_1(I_S)$ is equal to $\dim Gr(\mathbb{P}^{k-1}, \mathbb{P}^n) - \dim S - \dim \phi_2^{-1}(S) = k(n+1-k) - n + 2 - k(n-2-k)$. That calculation is effective thanks to the result of Chiantini and Ciliberto on the non-defectivity of the Grassmannians of secant varieties of curves (see [2]). We have that this variety has codimension $3k - n + 1$, but we are in the hypothesis $3k < n - 1$, so $3k - n + 1 < 0$.

For $3k = n - 1$ the condition gives $\mathrm{codim} = 0$, so it is verified for all L.

5. Restricted Tangent Bundle

For the restricted tangent bundle we obtain similar results as for the normal bundle.

Lemma 5.1. *Let $n \leq 2k$. The codimension in $Gr(\mathbb{P}^{k-1}, \mathbb{P}^n)$ of the variety of $L \cong \mathbb{P}^{k-1}$ in $\mathbb{P}^n$ belonging to some $(n-r)$-secant $\mathbb{P}^{n-r-1}$ to the rational normal curve in $\mathbb{P}^n$ is $k - n + r + kr$.*

In particular for $r = n - k - 1$ we have:

Theorem 5.2. *Let $n \leq 2k$. If the centre of projection $L \cong \mathbb{P}^{k-1}$ belongs to some $(k+2)-$secant $\mathbb{P}^{k+1}$ to the rational normal curve C_n in $\mathbb{P}^n$, then we have:*

$$T\mathbb{P}^{n-k}|_{\pi_k(C_n)} \cong \mathcal{O}(n+1)^{n-k-1} \oplus \mathcal{O}(n+1+k).$$

Theorem 5.3. *Let $2k \leq n$. If the centre of projection $L \cong \mathbb{P}^{k-1}$ belongs to some $\mathbb{P}^k$ $(k+1)-$secant to the rational normal curve C_n in $\mathbb{P}^n$, then we have:*

$$T\mathbb{P}^{n-k}|_{\pi_k(C_n)} \cong \mathcal{O}(n+1)^{n-k-1} \oplus \mathcal{O}(n+1+k).$$

Observation 5.4. *By Theorem 1 in* [10] *(see also* [12]*)* $T^n_{n-k}((n+1)^{n-k-1}, n+1+k)$ *is an irreducible variety of* $\operatorname{codim}(T^n_{n-k}((n+1)^{n-k-1}, n+1+k)) = (k-1)(n-k-1) \leq kn - k^2 - k - 1$.

Corollary 5.5. *Let $2k < n$. The variety of linear spaces $L \cong \mathbb{P}^{k-1}$ such that, $\pi_L(C_n) \subset \mathbb{P}^{n-k}$ has the restricted tangent bundle $T\mathbb{P}^{n-k}_{\pi_k(C_n),\mathbb{P}^{n-k}} \cong \mathcal{O}(n+1)^{n-k-1} \oplus \mathcal{O}(n+1+k)$ has an irreducible subvariety of codimension $kn - k^2 - k - 1$ in $Gr(\mathbb{P}^{k-1}, \mathbb{P}^n)$ formed by the linear spaces L belonging to some $(k+1)-$secant $\mathbb{P}^k$.*

ACKNOWLEDGEMENTS

This paper is part of my PhD thesis. I am very grateful to my advisor Professor Giorgio Ottaviani for the patience with which he followed this work very closely.

REFERENCES

[1] Alessandro Bernardi. Normal bundle of rational curves and waring decomposition. Arxiv e-print, March 2012.

[2] L. Chiantini and C. Ciliberto. The Grassmannians of secant varieties of curves are not defective. *Indag. Math. (N.S.)*, 13(1):23–28, 2002.

[3] D. Eisenbud and A. Van de Ven. On the normal bundles of smooth rational space curves. *Math. Ann.*, 256(4):453–463, 1981.

[4] David Eisenbud and A. Van de Ven. On the variety of smooth rational space curves with given degree and normal bundle. *Invent. Math.*, 67(1):89–100, 1982.

[5] Franco Ghione. Quelques exemples de courbes de $\mathbf{P}^3$ dont le fibré normal ne se décompose pas. *C. R. Acad. Sci. Paris Sér. A-B*, 285(5):A375–A377, 1977.

[6] Franco Ghione and Gianni Sacchiero. Normal bundles of rational curves in $\mathbf{P}^3$. *Manuscripta Math.*, 33(2):111–128, 1980.

[7] A. Gimigliano, B. Harbourne, and M. Idà. On plane rational curves and the splitting of the tangent bundle. *Accepted on Annali della Scuola Normale Superiore di Pisa, Classe di Scienze.*

[8] Anthony Iarrobino and Vassil Kanev. *Power sums, Gorenstein algebras, and determinantal loci*, volume 1721 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1999. Appendix C by Iarrobino and Steven L. Kleiman.

[9] Josep M. Miret. On the variety of rational curves in $\mathbf{P}^n$. *Ann. Univ. Ferrara Sez. VII (N.S.)*, 32:55–65 (1987), 1986.

[10] Luciana Ramella. La stratification du schéma de Hilbert des courbes rationnelles de $\mathbf{P}^n$ par le fibré tangent restreint. *C. R. Acad. Sci. Paris Sér. I Math.*, 311(3):181–184, 1990.

[11] Luciana Ramella. Sur les schémas définissant les courbes rationnelles lisses de $\mathbf{P}^3$ ayant fibré normal et fibré tangent restreint fixés. *Mém. Soc. Math. France (N.S.)*, 54:ii+74, 1993.

[12] J. Verdier. Two dimensional σ-models and harmonic maps from S^2 to S^{2n}. In M. Serdaroglu and E. Ínönü, editors, *Group Theoretical Methods in Physics*, volume 180 of *Lecture Notes in Physics*, pages 136–141. Springer Berlin / Heidelberg, 1983.

DIPARTIMENTO DI MATEMATICA, UNIVERSITY OF FLORENCE
E-mail address: `bernardi@math.unifi.it`